

Creation and Decay of η -Mesic Nuclei

G.A. Sokol¹, T.A. Aibergenov, A.V. Koltsov, A.V. Kravtsov,
Yu.I. Krutov, A.I. L'vov, L.N. Pavlyuchenko,
V.P. Pavlyuchenko, S.S. Sidorin

P.N. Lebedev Physical Institute, Leninsky Prospect 53, Moscow 117924, Russia

Abstract.

First experimental results on photoproduction of η -mesic nuclei are analyzed. In an experiment performed at the 1 GeV electron synchrotron of the Lebedev Physical Institute, correlated π^+n pairs arising from the reaction

$$\gamma + {}^{12}\text{C} \rightarrow N + {}_{\eta}(A-1) \rightarrow N + \pi^+ + n + (A-2)$$

and flying transversely to the photon beam have been observed. When the photon energy exceeds the η -meson production threshold, a distribution of the π^+n pairs over their total energy is found to have a peak in the subthreshold region of the internal-conversion process $\eta p \rightarrow \pi^+n$ which signals about formation of η -mesic nuclei.

The idea that a bound state of the η -meson and a nucleus (the so-called η -mesic nucleus) can exist in Nature was put forward long ago by Peng [1] who relied on the first estimates of the ηN scattering length $a_{\eta N}$ obtained by Bhalerao and Liu [2]. Owing to $\text{Re } a_{\eta N} > 0$, an average attractive potential exists between slow η and nucleons. This can result in binding ηA systems, provided the life time of η in nuclei is long enough [3]. Modern calculations [4,5] predict a rather strong ηN attraction which is sufficient for binding η in all nuclei with $A \geq 4$.

The very first experiments on searching for the η -mesic nuclei performed at BNL [6] and LAMPF [7] gave negative results. Meantime, studies of the reactions $p(d, {}^3\text{He})\eta$ [8,9], ${}^{18}\text{O}(\pi^+, \pi^-){}^{18}\text{Ne}$ [10], and $\vec{d}(d, {}^4\text{He})\eta$ [11] suggest that a quasi-bound ηA state is formed in these reactions [12,13].

In the present work we report on first results concerning formation of η -mesic nuclei in photoreactions. A very efficient trigger for searching for η -mesic nuclei [15] consists in detecting decay products of the η -mesic nucleus, viz. a πN pair produced in the reaction $\eta N \rightarrow \pi N$ inside the nucleus. Here η itself is produced at an earlier stage, in the reaction $\gamma N \rightarrow \eta N$ in our case. Both these reactions are mediated by the $S_{11}(1535)$ resonance which affects also a propagation of the

¹⁾ Talk given at CIPANP-2000 (May 22-28, 2000, Quebec). E-mail: gsokol@x4u.lebedev.ru

intermediate η in the medium (via multiple ηN rescattering) and leads to capturing slow η into a bound state (Fig. 1). Formation of the bound state of the η and the nucleus becomes possible when the momentum of the produced η is small (typically less than 150 MeV/c). This requirement suggests photon energies $E_\gamma = 650\text{--}850$ MeV as most suitable for creating η -nuclei.




FIGURE 1. Mechanism of creation and decay of η -mesic nuclei.

πN pairs emerging from η -mesic nucleus decays have an opening angle $\langle\theta_{\pi N}\rangle = 180^\circ$ and specific kinetic energies of their components (though smeared by the Fermi motion), $\langle E_\pi \rangle \simeq 300$ MeV, $\langle E_n \rangle \simeq 100$ MeV. Among four possible isotopic combinations π^+n , π^-p , π^0n , π^0p the first one is quite suitable for measuring energies of the particles.

Accordingly, in an experiment performed at the 1 GeV electron synchrotron of the Lebedev Physical Institute, correlated π^+n pairs arising from the reaction

$$\gamma + {}^{12}\text{C} \rightarrow N + \eta(A-1) \rightarrow N + \pi^+ + n + (A-2) \quad (1)$$

have been searched for. An experimental setup (Fig. 2) consisted of a carbon target $\varnothing 4\text{ cm} \times 4\text{ cm}$ and two time-of-flight scintillator spectrometers having a time resolution of $\delta\tau \simeq 0.1$ ns. A plastic anticounter A of charged particles (of the 90% efficiency), placed in front of the neutron detectors, and dE/dx layers, placed between start and stop detectors in the pion spectrometer, were used for a better identification of particles.




FIGURE 2. Layout of the experimental setup. Shown also time-of-flight spectra in the π (left) and n (right) spectrometers.

Strategy of measurements was as follows. Two bremsstrahlung-beam energies were used, $E_{\gamma\text{max}} = 650$ MeV and 850 MeV, i.e., well below and well above η

production threshold on the free nucleon which is 707 MeV. The first, “calibration” run was performed at 650 MeV with the spectrometers positioned at angles $\theta_n = \theta_\pi = 50^\circ$ around the beam. In that run, this was a quasi-free photoproduction $\gamma p \rightarrow \pi^+ n$ which dominated the observed yield of the $\pi^+ n$ pairs. Then, at the same “low” energy 650 MeV, the spectrometers were positioned at $\theta_n = \theta_\pi = 90^\circ$ (the “background” run). In such a kinematics, the quasi-free production did not contribute and the observed count was presumably dominated by double-pion production. At last, the third run (the “effect+background” run) was performed at the same $90^\circ/90^\circ$ position, however with the higher photon beam energy of 850 MeV, at which η mesons are produced too.




FIGURE 3. Raw $\pi^+ n$ event distributions over the particle velocities β_n and β_π for the “calibration” run (a), “background” run (b), and “effect+background” run (c).

In accordance with measured velocities of particles detected by the spectrometers, all candidates to the $\pi^+ n$ events were separated into three classes: fast-fast (FF), fast-slow (FS), and slow-slow (SS) events. The FF events mostly correspond to $\pi^0 \pi^0$ production which results in hitting detectors by photons or e^+/e^- . The FS events mostly emerge from the $\pi^+ n$ pairs. Comparing yields and time spectra in these runs (and, in particular, using the SS events for extrapolating and subtracting a background), we have found a clear excess of the FS events which appeared when the photon energy exceeded η production threshold. The total cross section

of photoproduction of such excess pairs, averaged over the photon energy range 650–850 MeV, was found to be about $\sigma_{\text{tot}}(\pi^+n) \simeq 10 \mu\text{b}$. See Ref. [14] for more details. In the present work a further analysis of the excess FS events is done and their energy characteristics are determined.




FIGURE 4. Corrected two-dimensional distributions over the velocities β of the π^+n events with the end-point energy of the bremsstrahlung spectrum $E_{\gamma\text{max}} = 850$ and 650 MeV.

In order to find kinetic energies of the neutron and pion, the velocities $\beta_i = L_i/ct_i$ of both the particles have to be determined. They are subject to fluctuations stemming from errors δt_i and δL_i in the time-of-flight t_i and the flight base L_i . Such fluctuations are clearly seen in the case of the ultra-relativistic FF events which have experimentally observed velocities close but not equal to 1 (see Fig. 3). Therefore, an experimental β -resolution of the setup can be directly inferred from the FF events. Then, using this information and applying an inverse-problem statistical method described in Ref. [16], one can unfold the experimental spectrum, obtain a smooth velocity distribution in the physical region $\beta_i < 1$ (Fig. 4), and eventually find a distribution of the particle's kinetic energies $E_i = M_i[(1 - \beta_i^2)^{-1/2} - 1]$. Finding E_i , we introduced corrections related with average energy losses of particles in absorbers and in the detector matter. It is worth to say that the number of the π^+n FS events visibly increases when the photon beam energy becomes sufficient for producing η mesons.

Of the most interest is the distribution of the π^+n events over their total energy $E_{\text{tot}} = E_n + E_\pi$, because creation and decay of η -mesic nuclei is expected to produce a relatively narrow peak in E_{tot} of the width ~ 50 – 70 MeV (see, e.g., [14,17]). Such a peak was indeed observed: see Fig. 5, in which an excess of the FS events appears when the photon energy exceeds the η -production threshold. Subtracting a smooth background, we have found a 1-dimensional energy distribution of the π^+n events presumably coming from (bound) η decaying in the nucleus, see Fig. 6. The experimental width of this distribution is about 100 MeV, including the apparatus resolution. Its center lies by $\Delta E = 40$ MeV below the energy excess $m_\eta - m_\pi = 408$ MeV in the reaction $\eta N \rightarrow \pi N$, and it is well below the position of the $S_{11}(1535)$ resonance too. Up to effects of binding of protons annihilated in




FIGURE 5. Distribution over the total kinetic energy of the π^+n pairs for the “effect+background” run (the left panel) and for the “background” run (the right panel) obtained after unfolding the raw spectra.




FIGURE 6. Distribution over the total kinetic energy of the π^+n pairs after subtraction of the background. Arrows indicate threshold in the reaction $\eta N \rightarrow \pi N$, i.e. 408 MeV, and the weighted center of the histogram. For a comparison, a product of free-particle cross sections of $\gamma N \rightarrow \eta N$ and $\eta N \rightarrow \pi N$ [5] is shown with the dashed line (in arbitrary units).

the decay subprocess $\eta p \rightarrow \pi^+n$, the value ΔE characterizes the binding energy of η in the nucleus. The width of that peak is determined both by the width of the η -bound state and by the Fermi motion.

Whereas the fixed opening angle $\theta_{\pi n} = 180^\circ$ chosen in the kinematics with $\theta_n = \theta_\pi = 90^\circ$ selects π^+n pairs carrying a low total momentum in the direction of the photon beam, an independent check of the transverse momentum $p_\perp = p_\pi - p_n$ is meaningful. The corresponding distribution is shown in Fig. 7. On the top of a background, there is a narrower peak in $p_\perp$ having a width compatible with the Fermi momentum of nucleons in the nucleus.

In conclusion, an excess of correlated π^+n pairs with the opening angle $\langle \theta_{\pi N} \rangle = 180^\circ$ has been experimentally observed when the energy of photons exceeded the η -production threshold. A distribution of the pairs over their total kinetic energy was found to have a peak lying below threshold of the elementary process $\pi N \rightarrow \eta N$.




FIGURE 7. Distribution over the total transverse momentum $p_{\perp}$ of π^+n pairs for the “effect+background” run (the left panel) and for the “calibration” run (the right panel).

A narrow peak is also found in the pair’s distribution over their total transverse momentum. All that suggests that these π^+n pairs arise from creation and decay of captured bound η in the nucleus, i.e., they arise through the stage of formation of an η -mesic nucleus.

The work was supported by RFBR grant 99-02-18224.

REFERENCES

1. J.C. Peng, AIP Conference Proceedings **133**, 255 (1985).
2. R.S. Bhalerao and L.C. Liu, *Phys.Rev.Lett.* **54**, 865 (1985).
3. L.C. Liu and Q. Haider, *Phys. Rev. C* **34** (1986) 1845.
4. M. Batinić et al., *Phys. Rev. C* **57** (1998) 1004.
5. A.M. Green and S. Wycech, *Phys. Rev. C* **55** (1997) R2167.
6. R.E. Chrien et al., *Phys. Rev. Lett.* **60** (1988) 2595.
7. B.J. Lieb and L.C. Liu, LAMPF Progress Report, LA-11670-PR (1988).
8. J. Berger et al., *Phys. Rev. Lett.* **61** (1988) 919.
9. B. Mayer et al., *Phys. Rev. C* **53** (1996) 2068.
10. J.D. Johnson et al., *Phys. Rev. C* **47** (1993) 2571.
11. N. Willis et al., *Phys. Lett. B* **406** (1997) 14.
12. C. Wilkin, *Phys. Rev. C* **47** (1993) R938.
13. L. Kondratyuk et al., Proc. Int. Conf. “Mesons and Nuclei at Intermediate Energies”, Dubna 1994 (World Scientific, Singapore, Eds. M.Kh. Khanhasayev and Zh.B. Kurmanov), p. 714.
14. G.A. Sokol et al., *Fizika B* **8** (1998) 81.
15. G.A. Sokol and V.A. Tryasuchev, *Kratk. Soobsh. Fiz.* [Sov. Phys. – Lebedev Institute Reports] **4** (1991) 23.
16. V.P. Pavlyuchenko, *Vopr. Atom. Nauki i Tekhn, ser. Tekh. Fiz. Eksp.* **1/13** (1983) 39.
17. A.I. L’vov, nucl-th/9810054 and Proc. 7th Int. Conf. “Mesons and Light Nuclei”, Pruhonice, Prague, Czech Republic, 1998 (World Scientific, eds. J. Adam et al.), p. 469.